

1 Full title: **Estimating day range from camera-trap data:**  
2 **the animals' behaviour as a key parameter**

3 Running title: Estimating day range from camera-traps

4 **Pablo Palencia<sup>a\*</sup>, Joaquín Vicente<sup>a</sup>, Patricia Barroso<sup>a</sup>, José Ángel Barasona<sup>a,b</sup>,**  
5 **Ramón C. Soriguer<sup>c</sup> & Pelayo Acevedo<sup>a</sup>**

6 <sup>a</sup>Instituto de Investigación en Recursos Cinegéticos, IREC (UCLM-CSIC-JCCM), C/  
7 Ronda de Toledo 12, 13071 Ciudad Real, Spain.

8 <sup>b</sup>VISAVET Centre, Animal Health Department, Universidad Complutense de Madrid,  
9 Avd. Puerta de Hierro, s/n, Facultad de Veterinaria, 28040, Madrid, Spain;

10 <sup>c</sup>Estación Biológica de Doñana (EBD-CSIC), C/ Américo Vespucio, s/n, 41092, Sevilla,  
11 Spain.

12 \*Corresponding author: Instituto de Investigación en Recursos Cinegéticos, IREC  
13 (UCLM-CSIC-JCCM), Ronda de Toledo 12, 13071, Ciudad Real, Spain. Tel: 34  
14 926295300. E-mail. [pablo.palencia@uclm.es](mailto:pablo.palencia@uclm.es)

## Abstract

Day Range (DR), the distance travelled by an individual during the day, is an important metric in movement ecology. Moreover, DR is an important parameter for estimating population density through Random Encounter Model (REM). Traditionally, DR has been estimated applying GPS technology and considering straight-line distances between consecutive locations, which suppose an underestimation of the true path distance. In this work, we have tested the accuracy of a new approach based on camera-trap data for the estimation of DR when including animals' behaviour in the formulation. For that, we have considered wild boar (*Sus scrofa*) as model species. We tagged 18 individuals with telemetry devices and then monitored the population with camera-traps (photo and video mode) in order to estimate the DR. In the case of telemetry, a straight-line DR was estimated and rescaled with a tortuosity-related correction factor. Using this camera-trap data, we have revisited the procedure described by Rowcliffe *et al.* (2016) to estimate the DR. A new derivation of this approach was then developed, in which different animal behaviours were weighted to estimate the DR. There were no significant differences between the DR values obtained using telemetry data (corrected by the tortuosity-related correction factor) and those attained with the new approach. However, the original approach employed to estimate the DR based on camera-trap data underestimated this parameter. The DR estimated with the weighted approach is  $12.74 \text{ Km} \cdot \text{day}^{-1} \pm (\text{SE}) 1.89$ . Here, we showed that animals' behaviour should be weighted in order to estimate the DR when working with species that behave differently in front of the cameras. Our results described a new approach for estimating movement parameters without the need of capture and tagged individuals. These results can have a lot of relevance in movement ecology, disease dynamics, and population monitoring methods.

**Keywords:** Animal movement, population monitoring, Random Encounter Model, telemetry, tortuosity, wild boar.

## 1. Introduction

Obtaining accurate estimates of population density continues to be a constant challenge for wildlife management and conservation (Nichols & Williams, 2006). It is widely recognized that obtaining population size estimations is costly and that their reliability is usually not sufficient for management purposes, being necessary to consider other parameters related to body condition, vegetation effects, etc. (Morellet *et al.*, 2007). Some species are currently undergoing a notable expansion worldwide (e.g. wild boar *Sus scrofa*; Massei *et al.*, 2015), whereas others are substantially declining in their entire distribution range (e.g. farmland birds; Inger *et al.*, 2015). Feasible methods with which to attain precise and accurate estimations of population density are, therefore, in great demand in both scenarios. In this respect, the use of remotely triggered cameras (camera-traps, CT) for this purpose has substantially increased over the last few years (Rovero & Zimmermann, 2016). Even camera-traps can be detected by wildlife (Meek *et al.*, 2014) is a method less intrusive than others, and most importantly, makes it possible to obtain information on highly cryptic species inhabiting a wide range of habitats (Steenweg *et al.*, 2017). CT are, therefore, now firmly established as a core tool for wildlife monitoring.

CT were traditionally used in wildlife monitoring in order to derive relative indices of population abundance (e.g. Massei *et al.*, 2017). When population densities are the target parameter, CT are used to obtain data for capture-recapture methods (CR) and, more recently, spatially explicit approaches have been developed (Royle & Young, 2008; Jiménez *et al.*, 2017). However, the use of CR models (in a broad sense) requires marking individuals or working with individually recognisable species (but see spatial counts; Chandler & Royle, 2013). The use of these methods requires a greater effort than that of others. This is because it is necessary to make individuals recognisable for CT, since only a few species have a sufficient variability of natural marks for them to be individually recognisable. Rowcliffe *et al.* (2008) developed a method with which to estimate animal density using CT without individual recognition, the Random Encounter

Model (REM). The REM describes the rate of contact between animals and CT to estimate population density. For that requires trapping rate (i.e. the number of passages per unit of time), the day range (DR, i.e. the distance travelled by an individual in a day), mean group size and the radius and angle of the camera's field of detection. Although it is not as easy to estimate camera-related parameters as it is to consult model specifications (Rowcliffe *et al.*, 2011; Hofmeester, Rowcliffe & Jansen, 2017), the DR is the most costly and time-consuming parameter required for REM application (Rowcliffe *et al.*, 2012; Prichard *et al.*, 2014; Alexander & Maritz, 2015; Sennhenn-Reulen *et al.*, 2017; Nakashima, Fukasawa & Samejima, 2018). This signifies that, despite being a promising method for implementation in wildlife monitoring programmes, even on large spatial scales (Manzo *et al.*, 2012; Zero *et al.*, 2013; Anile *et al.*, 2014; Cusack *et al.*, 2015; Balestrieri *et al.*, 2016; Rademaker *et al.*, 2016; Caravaggi *et al.*, 2017; Gray, 2018), the application of REM is, at present, considerably limited by the estimation of the DR (Rovero & Marshall, 2009; Alexander & Maritz, 2015; Cusack *et al.*, 2015; Pfeffer *et al.*, 2017). Recently, Nakashima *et al.* (2018) described the Random Encounter and Staying Time (REST), an extension of the REM based on the amount of time that detected animals remain in the field of view of the CT rather than the DR; but no field trails have yet been undertaken to test the applicability of this method.

The DR has traditionally been estimated by employing intermittent location data, usually derived from telemetry studies (mainly using GPS devices), and then adding up the straight-line distances between consecutive locations in order to attain the total distance (Sennhenn-Reulen *et al.*, 2017; Tucker *et al.*, 2018). However, this approach underestimates the DR because the actual paths between consecutive relocations are not, in fact, straight (Laundre *et al.*, 1987; Reynolds & Laundre, 1990; Rowcliffe *et al.*, 2012; Sennhenn-Reulen *et al.*, 2017); animal paths are mostly tortuous (i.e. characterised by continuous turns). Estimates obtained from low sampling frequencies are, therefore, significantly shorter than those based on higher sampling frequencies. In order to manage these tortuosity-related problems, some authors have used more

intensive monitoring programmes with a higher frequency of relocations of tagged animals. This does not, however, solve the problem of tortuosity and other problems emerge, which are mainly related to fixed accuracy and battery life (Johnson & Ganskopp, 2008; Hurford, 2009; Dewhirst *et al.*, 2016); it is not really possible to register a complete track of tagged individuals with the current telemetry devices Dewhirst *et al.*, 2016, Sennhenn-Reulen *et al.*, 2017). Some authors have proposed tracking animals in the snow or the visual tracking of wild animals in parallel with a continuous GPS location series (Musiani, Okarma & Jędrzejewski, 1998; Sennhenn-Reulen *et al.*, 2017), but these approaches are limited to certain specific areas and species that are accustomed to human observers. Moreover, this approach is costly and time-consuming. In this context, various studies have verified correction factors with which to rescale the values obtained from intermittent fixed data to precise values of the DR and their applicability (Pépin *et al.* 2004; Rowcliffe *et al.*, 2012). Nevertheless, Rowcliffe *et al.* (2016) recently developed an approach with which to estimate the DR based solely on the information derived from CT. In this approach, the DR is estimated as the product of travel speed (i.e. average speed while active) and activity level (i.e. the proportion of the day that the population is active).

In this work we have considered the wild boar as a model species. The wild boar is widely distributed throughout Europe, with a high ecological and social relevance (Acevedo *et al.*, 2014; Massei *et al.*, 2015), particularly owing to its role as a reservoir of relevant shared diseases like tuberculosis or African Swine Fever (Gortazar *et al.*, 2012; Blome, Gabriel & Beer, 2013; Keuling *et al.*, 2018). Reliable estimates of population densities are needed for further risk assessment, essential to improve management strategies (Keuling *et al.*, 2018). However, the methods traditionally employed to estimate wildlife population density are not reliable when working with this species owing to its behavioural peculiarities (but see Engeman *et al.*, 2013). The nocturnal habits and elusive behaviour of the wild boar limited the applicability of those methods based on direct observation; and indirect methods, like pellets counts, are limited by the

estimations of other local-parameters such as defecation rates (Keuling *et al.*, 2018). For all of the above, the REM can be a potentially feasible option (Chauvenet *et al.*, 2017) once the method has been optimized, mostly as regards estimating the DR from CT data. Moreover, movement-related parameters can be essential for understanding and optimizing disease control efforts (Podgórski & Śmietanka, 2018)

While different approaches to derive DR values have been proposed, comparative studies in which the precision and accuracy of the estimations have been tested under field conditions are scarce. The aim of this study is to test a new approach for the estimation of the DR from CT using different categories of behaviour: feeding and moving. We hypothesized that for those species with marked behaviour could be necessary to weight the speed and activity values to finally estimate a DR value. Moreover, we are going to compare the DR estimations obtained with CT in photo and video mode. At this respect, we hypothesized that the estimations obtained in photo mode (using rapid-fire technology) can be similar to those obtained in video mode. These result could enhance the use of photo mode because of the advantages in relation to video (Glen *et al.*, 2013; Rovero & Zimmermann, 2016; Steenweg *et al.*, 2017).

## **2. Materials and methods**

### **2.1. Study area**

The study was carried out in the Doñana Biological Reserve (DBR), a territory of approximately 6800 ha located in Doñana National Park (DNP) (37°0'N, 6°30'W). DNP is a natural area covering 54000 ha located on the Atlantic coast of southwest Spain. As it is a protected area, neither hunting activities nor any management actions, such as artificial feeding or the provision of water for wild ungulates, are permitted.

### **2.2. Day range estimation**

Eighteen wild boars were tagged with GPS-GPRS (Microsensory System, Sevilla, Spain) collars in three different periods to estimate their DR. The population was also monitored with CT during one of the telemetry study periods with the same purpose.

#### 2.2.1. Telemetry

The animals were captured in different trapping areas throughout the DBR using five padded foothold cage traps in order to collar a sample of animals from multiple social groups. The animals were captured and managed following a protocol approved by the Animal Experiment Committee of Castilla-La Mancha University and by the Spanish Ethics Committee (PR-2015-03-08), designed and developed by scientists (B and C animal experimentation categories) in accordance with EC Directive 86/609/EEC for animal handling and experiments. The wild boar was anaesthetised following the protocol described by Barasona *et al.* (2013), weighed, ear tagged, collared with GPS-GPRS devices and assessed for body condition, age and sex. In those cases that more than one individual was captured at the same time, we only collared one of them to avoid pseudoreplication problems in telemetry data. The positional error associated with the GPS locations was 12.08 m (SE = 0.20 m). The data collected included the date, time, geographic coordinates and location acquisition time (LAT, ranges between 0–255 s). Those GPS locations with LAT = 255 s were screened to detect and then remove anomalous fixes (manufacturer's technical data). The locations obtained during the days of collar deployment and of collar retrieval were discarded for the analysis to avoid possible anomalous behaviour associated with handling procedures. .

Three independent study periods and sampling intensities have been considered in this work. Nine adults (>24 months) wild boar (1 female and 8 males) were tagged between December 2012 - March 2013 (period 1); with collars programmed to acquire one location every hour (i.e. 24 locations·day<sup>-1</sup>). Four adults wild boar (2 female and 2 males) were tagged between December 2015 - March 2016 (period 2); with collars programmed to acquire one location every two hours (i.e. 12 locations·day<sup>-1</sup>). Five adults wild boar (3 females and 2 males) were tagged between January 2017 - March 2017

(period 3); with collars programmed to acquire one location every five minutes (i.e. 288 locations·day<sup>-1</sup>). Inter-sex significant differences were not found in the movement rate of the tagged individuals (linear model, Period 1-p:0.362; Period 2-p:0.655) and all individuals were, therefore, pooled in the analyses (see below). The collars were removed when animals died or were shot while the population control programme was being carried out.

### 2.2.2. Camera-traps

The CT survey was carried out between January-February 2017 (period 3). We designed a 1.5Km grid to guide the location of the 16 sampling points throughout the DBR, covering all the habitats in the study area (Rowcliffe *et al.*, 2013) (Fig. 1). Two Little Acorn cameras (Ltl-5310 Series LED IR Invisible) were placed at each point, without bait (Rowcliffe *et al.*, 2016), one of them in photo mode and the other in video mode. The CT were located within a buffer of 100m around the nodes of the grid in trees or man-made poles. The motion sensitivity of all the cameras was set to high. The cameras were deployed on the tree nearest to the computer-generated point, 30-50 cm above the ground, with the angle of view parallel to the overall slope and unobstructed by vegetation. If necessary, the vegetation in front of the camera was cut to reduce false triggers resulting from its movement. The cameras were set to be operative all day, with the use of an infrared flash at night, and the date and time of capture were automatically stamped onto each image. The cameras were triggered by means of passive infrared motion sensors and recorded a sequence of three consecutive photos or 30s-long video clips. In order to record the individuals' trajectories in as much detail as possible, the minimum time lapse between bursts was selected (1 second). The cameras were checked every 10 days in order to change their batteries and SD cards, and to ensure that they were functioning correctly.

Six markers (wooden sticks of 1m in length) were placed in front of the camera forming an arc covering the angle of vision of the camera, three of them at 5m from the



camera, and the other three at 10m from the camera (Caravaggi *et al.*, 2016; Hofmeester *et al.*, 2017; see Fig. 2). After taking one photo of the structure, the sticks were removed. These marks were later used to locate the individuals captured with the cameras with a precision of 0.5m, and to estimate their travel speed. We have considered only those passages below a distance of 10m from the camera.

### 2.2.3. Approaches to derive DR

#### *Telemetry: approach-1*

The daily movement between consecutive locations was calculated using a programmed function in R 3.3.3 (Appendix 1). This was done by calculating the sum of the straight-line distances between consecutive fixes.

For periods 1 and 2, we have considered only those days with more than 80% of the fixed rate and a daily movement value was estimated for each day. For period 3, an average movement speed ( $\text{m}\cdot\text{s}^{-1}$ ) was calculated for each hour of the day in order to subsequently obtain the daily mean speed. The daily speed ( $\text{m}\cdot\text{s}^{-1}$ ) was then directly transformed into the DR ( $\text{Km}\cdot\text{day}^{-1}$ ). This analytical procedure of pooling data per hour was adopted owing to the small number of days on which collars captured good fixes for the entire daily range. In the three periods, the DR values obtained directly from telemetry data were corrected using the tortuosity-related correction factor developed by Rowcliffe *et al.* (2012). The correction factor was calculated as the inverse value of the accuracy obtained, with the objective of rescaling the DR directly obtained from telemetry data to more accurate DR estimations. This correction was shown to be similar between species (see Fig.4 in Rowcliffe *et al.*, 2012) with a relation between the body mass and the underestimation at a given sampling frequency. The correction factor increases as the sampling intensity decrease. For our data, we have considered the correction factor estimated for peccary (*Tayassu tajacu*), because this species is similar to wild boar in biological terms and as regards its body mass.

### Camera-traps: approach-2 & approach-3

The procedure by which to estimate the DR from information obtained from CT described by Rowcliffe *et al.*, (2016) was then applied. In this method, travel speed is estimated using the photos/videos of the animals captured by the CT. This is done by dividing the distance travelled by the duration of the sequence (difference in time between the time stamps on the first and last picture/photogram). As suggested by Rowcliffe *et al.* (2016), those sequences in which animals reacted to the camera were discarded. However, animals are more likely to make contact with cameras when they move faster (Hutchinson & Waser, 2007). This problem was solved by fitting log-normal and Weibull distribution models in order to estimate the average travel speed (Rowcliffe *et al.*, 2016); the most parsimonious models were selected using AIC.

Camera-traps only detect animals when they are outside their refuges (i.e. active animals) and it is, therefore, necessary to take into account the population's activity rate. The data obtained from the CT were also employed to estimate activity (Rowcliffe *et al.*, 2014). This procedure is based on the key assumption that all the individuals in the population being sampled are active at the peak of the trap rate, and the trap rate at a given time of day is, therefore, proportional to the population's level of activity. The activity index per day was estimated using the 'activity' R package (Rowcliffe, 2016). In approach-2, the DR was, therefore, estimated as the product of the activity index ( $a$ ) and averaged travel speed ( $s$ ).

However, wild boars were recorded by the CT with two marked different behaviours. The behaviours observed with the cameras can be classified in two categories with quite different movement speeds: i) feeding and, therefore, slow movements, and ii) moving among habitats and, therefore, rapid movements. The general activity value was multiplied by the percentage of sequences of each type of behaviour to obtain an activity value for each of them. This consequently led us to describe a new procedure, approach-3, in which independent activity indices ( $a_i$ ) and travel speeds ( $s_i$ ) are estimated for each behaviour and, therefore, in this case,  $DR = (a_{\text{feed}} \cdot s_{\text{feed}}) + (a_{\text{move}} \cdot s_{\text{move}})$ .

Approaches 2 and 3 were applied in this study using the data obtained from both photos (approaches 2.1 and 3.1, respectively) and videos (approaches 2.2 and 3.2, respectively). The standard error in these approaches was estimated by means of Goodman's (1960) variance of product formula.

The DR values obtained using the different approaches were then statistically compared using the Wald test, with a test statistic  $W$  assessed on the chi-squared distribution with one degree of freedom (Wald & Wolfowitz, 1940).

### 3. Results

#### 3.1. Approach-1

One hundred and twenty-eight daily movements were estimated for period 1, 132 for period 2, and 2743 travel speeds for period 3. The mean straight-line DR values for each period were:  $2.91 \text{ Km} \cdot \text{day}^{-1} \pm 0.24$ ,  $1.96 \text{ Km} \cdot \text{day}^{-1} \pm 0.17$ , and  $6.19 \text{ Km} \cdot \text{day}^{-1} \pm 0.03$ , respectively. After applying the tortuosity-related correction factor (6.40 for period 1, 8.78 for period 2, and 2.19 for period 3, Rowcliffe *et al.*, 2012), the following corrected DRs were obtained:  $18.62 \text{ Km} \cdot \text{day}^{-1} \pm 2.91$  for period 1,  $17.21 \text{ Km} \cdot \text{day}^{-1} \pm 2.24$  for period 2, and  $13.55 \text{ Km} \cdot \text{day}^{-1} \pm 0.56$  for period 3.

#### 3.3. Approaches 2 and 3

The activity indices and travel speeds estimated using the CT data are shown in Table 2. The activity patterns are shown in Fig 3. The average speed for moving behaviour is greater than that obtained for feeding (Table 2). In photo mode, 212 individuals were recorded in 149 passages. A DR of  $4.93 \text{ Km} \cdot \text{day}^{-1} \pm 1.02$  (approach-2.1) and a weighted DR of  $12.74 \text{ Km} \cdot \text{day}^{-1} \pm 1.89$  (approach-3.1) were obtained from these data. In video mode, 141 individuals were recorded in 102 passages. The video data produced a DR of  $3.01 \text{ Km} \cdot \text{day}^{-1} \pm 1.06$  (approach-2.2) and a weighted DR of  $14.00 \text{ Km} \cdot \text{day}^{-1} \pm 2.13$  (approach-3.2).

Pairwise comparisons of the DR estimates between approaches showed only statistically significant differences ( $p < 0.05$ ) as regards the DR values obtained from approaches 2.1 and 2.2 and the other approaches tested (Fig. 4).

#### 4. Discussion

We have revisited the approach by Rowcliffe *et al.* (2016) and we weighted by the behaviour the formula with which to estimate the DR based on CT data. We have done this by considering the results obtained in the same population monitored with telemetry devices as a reference value (after applying the tortuosity-related correction factor) for the DR. While the approach generally used to estimate the DR on the basis of CT data (approach-2) notably underestimated the DR, the approach weighted by the behaviours (approach-3) produced DR estimates that were coherent with those derived from telemetry data, after correction with the tortuosity factor.

The problem of underestimating the DR from telemetry data is well known (Rowcliffe *et al.*, 2012; Alexander & Maritz, 2015; Sennhenn-Reulen *et al.*, 2017) and have been evidenced again in this work. The DR values estimated through approaches 1 & 3 are around 15Km/day. Although these results are very different from those usually obtained for wild boar (e.g. Spitz & Janeau, 1990; Podgórski *et al.*, 2013), it can be explained based on the fact that most of these studies considered straight-line distances between consecutive locations obtained in telemetry studies. These studies, even it can be very useful to compare animal behaviour between different situations (e.g. Podgórski *et al.*, 2013), are limited in the accurate of the results obtained in relation with DR. To our knowledge, there are no studies that have been estimated the “real” DR of wild boar.

Camera-traps have been proposed as a useful method with which to estimate movement parameters in accordance with their higher spatial accuracy of a few centimeters (Rowcliffe *et al.*, 2012, 2016; Dewhurst *et al.*, 2016). In this paper, we have employed the data obtained from CT as a basis on which to obtain DR values equivalent to those obtained by means of telemetry. However, we have developed a new approach

based on that described by Rowcliffe *et al.* (2016). In this respect, the DR estimates obtained by directly applying the methodology described by Rowcliffe *et al.* (2016) – approach-2– were significantly lower than those obtained with the maximum sampling frequency used on tagged animals, even though frequencies such as those used in this work underestimate the real value by 50% (Rowcliffe *et al.*, 2012). This evidences that approach-2 is not able to derive precise DR values, or at least not with species that have marked behaviours. With these species, it is necessary to estimate a particular travel speed and activity for each particular type of behaviour, and then derive weighted DR estimations. In this study, after working with wild boar, we have proposed two general behaviours: “moving” and “feeding”. We will eventually be able to record other behaviour, such as that of rutting. Our classification is based on the precise speed for each type of behaviour, instead of ecological significance. The types of behaviour observed should, therefore, eventually be classified on the basis of their speed. Rapid movements should be included in the feeding group, and slow movements in the moving group.

Most of the species with which the original approach for estimating DR from CT data was previously implemented were carnivores and rodents (Rowcliffe *et al.*, 2016). Carnivore feeding behaviour can be described as a succession of feeding bouts and relatively straight and rapid relocation movements towards patches offering new feeding resources (Getz & Saltz, 2008). These species are not often recorded feeding, showing that most of the captures appertain to animals in movement. In the case of rodents, their movement have been described (Dobly, 2001; Cook, Anderson & Schweiger, 2004) as a succession of straight movements following natural or artificial linear structures (“runways”) to travel between zones with high availability of resources. This can be explained on the basis of the distribution of resources: while the distribution of carnivores’ or rodents’ food resources is very precise, ungulates have general and widely-distributed resources, and capturing individuals behaving in different manners is, therefore, more probable. For all of the above reasons, we propose that for those species that behave differently in front of the cameras, it might be necessary to apply the approach described

here in order to obtain precise estimations of the DR when using CT.

With regard to the estimates obtained from CT, no significant differences were found between photo and video modes. In this study, we have used the rapid-fire technology (i.e. time lapses between consecutive photos in the same series of less than one second) that allowed us to record the paths travelled by animals with great precision. We, therefore, suggest the use of photos rather than videos. This is principally because the trigger speeds and recovery times are generally longer in video mode when compared to photo mode (Rovero & Zimmermann, 2016; Nakashima *et al.*, 2018), and photo data can be processed more quickly than video data, which is one of the most time-consuming stages in CT studies (Steenweg *et al.*, 2017). But this is also because photos require considerably less memory than videos (Glen *et al.*, 2013). Moreover, the video data estimates were highly variable. This could be explained by the smaller sample size obtained. Even though photo and video cameras were deployed on the same tree/stick, they did not record the same passages. This could be explained by the fact that we have considered all passages below a distance of ten meters from the camera. Some authors who have evaluated the sensitivity of the camera have obtained an effective detection distance for wild boar of around 5-6 meters (Hofmeester *et al.*, 2017). These results lead us to conclude that those wild boar that passed between five and ten meters in front of the camera may not have been detected by both devices on all occasions.

Moreover, we have included a cost-benefit analyses (Appendix 2) comparing CT and telemetry as methods for estimating DR. Telemetry (always corrected by tortuosity) can be recommendable if we are working only with one species and we are interested also in other parameters like habitat use, CT have other advantages. For example, we are able to estimate DR for more than one species, and other ecology parameters like species occupancy and richness or activity patterns. Moreover, the applicability of CT studies are continually improving (e.g. Tabak *et al.*, 2018), and the method will be more efficient in the future. An especial situation where CT are highly recommendable is if we are applying the REM. In this case, the use of CT are mandatory to apply the REM, and

we can obtain all the parameters to estimate population density from the CT data, i.e. without the need from including other methods (e.g. telemetry).

Although, the variation in the movement behaviour of populations of the same species living in similar habitats could be minimal (Rowcliffe *et al.*, 2012), we recommend more studies for a better understanding of the variation in the DR between populations, habitats and seasons. Overall, the results reported here highlight the utility of the REM when estimating wildlife population density, since precise DR estimates can be derived from the CT and from the same survey used to estimate the trapping-rate or group size. But in this respect, animals' behaviour should be taken into account in order to derive precise measures of the DR. This study supposes a step forward as regards establishing the REM as a core tool for wild boar monitoring, for which the feasibility and reliability of traditional population monitoring methods are limited. Moreover, the movement behaviour of the species have a lot of relevance in other ecological studies (Morelle *et al.*, 2015), for example, as predictors of some diseases dynamics (Podgórski, Apollonio & Keuling, 2018).

## Acknowledgments

The authors would like to thank Dr. Marcus Rowcliffe for his continuous advice as regards estimating the DR from camera-trap data. They would also like to acknowledge Doñana National Park for providing the logistic support required to carry out this study. This work was partly funded by project AGL2016-76358-R, Ministerio de Economía, Industria y Competitividad (MINECO, FEDER-UE) and partly by an ENETWILD contract (<http://www.enetwild.com>). P.P. received supported from the MINECO-UCLM through an FPU grant (FPU16/00039) and from a UCLM grant during his Master's degree studies MUIBARC (<https://muibarc.masteruniversitario.uclm.es/presentacion.aspx>). P.B. received supported from ENETWILD. J.A.B. is supported by a Juan de la Cierva contract from MINECO-UCM (FJCI-2015-23643) and P.A. by MINECO-UCLM through a Ramón y Cajal contract (RYC-2012-11970).

## References

- Acevedo, P., Quirós-Fernández, F., Casal, J. & Vicente, J. (2014). Spatial distribution of wild boar population abundance: Basic information for spatial epidemiology and wildlife management. *Ecol. Indic.***36**, 594–600.
- Alexander, G.J. & Maritz, B. (2015). Sampling interval affects the estimation of movement parameters in four species of African snakes. *J. Zool.***297**, 309–318.
- Anile, S., Ragni, B., Randi, E., Mattucci, F. & Rovero, F. (2014). Wildcat population density on the Etna volcano, Italy: a comparison of density estimation methods. *J. Zool.***293**, 252–261.
- Balestrieri, A., Ruiz-Gonzalez, A., Vergara, M., Capelli, E., Tirozzi, P., Alfino, S., Minuti, G., Prigioni, C. & Saino, N. (2016). Pine marten density in lowland riparian woods: A test of the Random Encounter Model based on genetic data. *Mamm. Biol.***81**, 439–446.
- Barasona, J., López-Olvera, J., Beltrán-Beck, B., Gortázar, C. & Vicente, J. (2013). Trap-effectiveness and response to tiletamine-zolazepam and medetomidine anaesthesia in Eurasian wild boar captured with cage and corral traps. *BMC Vet. Res.***9**, 107.
- Blome, S., Gabriel, C. & Beer, M. (2013). Pathogenesis of African swine fever in domestic pigs and European wild boar. *Virus Res.***173**, 122–130.
- Caravaggi, A., Zaccaroni, M., Riga, F., Schai-Braun, S.C., Dick, J.T.A., Montgomery, W.I. & Reid, N. (2016). An invasive-native mammalian species replacement process captured by camera trap survey random encounter models. *Remote Sens. Ecol. Conserv.***2**, 45–58.
- Caravaggi, A., Banks, P.B., Burton, A.C., Finlay, C.M.V., Haswell, P.M., Hayward, M.W., Rowcliffe, M.J. & Wood, M.D. (2017). A review of camera trapping for conservation behaviour research. *Remote Sens. Ecol. Conserv.***3**, 109–122.
- Chandler, R.B. & Royle, J.A. (2013). Spatially explicit models for inference about density



439 in unmarked or partially marked populations. *Ann. Appl. Stat.***7**, 936–954.

440 Chauvenet, A.L.M., Gill, R.M.A., Smith, G.C., Ward, A.I. & Massei, G. (2017). Quantifying  
 441 the bias in density estimated from distance sampling and camera trapping of  
 442 unmarked individuals. *Ecol. Modell.***350**, 79–86.

443 Cook, W.M., Anderson, R.M. & Schweiger, E.W. (2004). Is the matrix really inhospitable?  
 444 Vole runway distribution in an experimentally fragmented landscape. *Oikos* **104**, 5–  
 445 14.

446 Cusack, J.J., Swanson, A., Coulson, T., Packer, C., Carbone, C., Dickman, A.J.,  
 447 Kosmala, M., Lintott, C. & Rowcliffe, J.M. (2015). Applying a random encounter  
 448 model to estimate lion density from camera traps in Serengeti National Park,  
 449 Tanzania. *J. Wildl. Manage.***79**, 1014–1021.

450 Dewhirst, O.P., Evans, H.K., Roskilly, K., Harvey, R.J., Hubel, T.Y. & Wilson, A.M.  
 451 (2016). Improving the accuracy of estimates of animal path and travel distance  
 452 using GPS drift-corrected dead reckoning. *Ecol. Evol.***6**, 6210–6222.

453 Doby, A. (2001). Movement patterns of male common voles (*Microtus arvalis*) in a  
 454 network of Y junctions: role of distant visual cues and scent marks. *Can. J. Zool.*  
 455 **79**, 2228–2238.

456 Engeman, R.M., Massei, G., Sage, M. & Gentle, M.N. (2013). Monitoring wild pig  
 457 populations: a review of methods. *Environ. Sci. Pollut. Res.***20**, 8077–8091.

458 Getz, W.M. & Saltz, D. (2008). A framework for generating and analyzing movement  
 459 paths on ecological landscapes. *Proc. Natl. Acad. Sci. U. S. A.***105**, 19066–71.

460 Glen, A.S., Cockburn, S., Nichols, M., Ekanayake, J. & Warburton, B. (2013). Optimising  
 461 Camera Traps for Monitoring Small Mammals. *PLoS One***8**, e67940.

462 Goodman, L.A. (1960). On the Exact Variance of Products. *J. Am. Stat. Assoc.***55**, 708.

463 Gortazar, C., Delahay, R.J., McDonald, R.A., Boadella, M., Wilson, G.J., Gavier-Widen,  
 464 D. & Acevedo, P. (2012). The status of tuberculosis in European wild mammals.  
 465 *Mamm. Rev.***42**, 193–206.

466 Gray, T.N.E. (2018). Monitoring tropical forest ungulates using camera-trap data. *J.*  
467 *Zool.***305**, 173–179.

468 Hofmeester, T.R., Rowcliffe, J.M. & Jansen, P.A. (2017). A simple method for estimating  
469 the effective detection distance of camera traps. *Remote Sens. Ecol. Conserv.***3**,  
470 81–89.

471 Hurford, A. (2009). GPS Measurement Error Gives Rise to Spurious 180° Turning Angles  
472 and Strong Directional Biases in Animal Movement Data. *PLoS One***4**, e5632.

473 Hutchinson, J.M.C. & Waser, P.M. (2007). Use, misuse and extensions of “ideal gas”  
474 models of animal encounter. *Biol. Rev.***82**, 335–359.

475 Inger, R., Gregory, R., Duffy, J.P., Stott, I., Voříšek, P. & Gaston, K.J. (2015). Common  
476 European birds are declining rapidly while less abundant species’ numbers are  
477 rising. *Ecol. Lett.***18**, 28–36.

478 Jiménez, J., Higuero, R., Charre-Medellin, J.F. & Acevedo, P. (2017). Spatial mark-  
479 resight models to estimate feral pig population density. *Hystrix Ital. J. Mammal.***28**,  
480 208–213.

481 Johnson, D.D. & Ganskopp, D.C. (2008). GPS collar sampling frequency: effects on  
482 measures of resource use. *Rangel. Ecol. Manag.***61**, 226.

483 Keuling, O., Sange, M., Acevedo, P., Podgorski, T., Smith, G., Scandura, M., Apollonio,  
484 M., Ferroglio, E. & Vicente, J. (2018). Guidance on estimation of wild boar  
485 population abundance and density: methods, challenges, possibilities. *EFSA*  
486 *Support. Publ.***15**.

487 Laundre, J.W., Reynolds, T.D., Knick, S.T. & Ball, I.J. (1987). Accuracy of Daily Point  
488 Relocations in Assessing Real Movement of Radio-Marked Animals. *J. Wildl.*  
489 *Manage.***51**, 937–940.

490 Manzo, E., Bartolommei, P., Rowcliffe, J.M. & Cozzolino, R. (2012). Estimation of  
491 population density of European pine marten in central Italy using camera trapping.  
492 *Acta Theriol. (Warsz.)***57**, 165–172.

493 Massei, G., Kindberg, J., Licoppe, A., Gačić, D., Šprem, N., Kamler, J., Baubet, E.,

- Hohmann, U., Monaco, A., Ozoliņš, J., Cellina, S., Podgórski, T., Fonseca, C., Markov, N., Pokorny, B., Rosell, C. & Náhlik, A. (2015). Wild boar populations up, numbers of hunters down? A review of trends and implications for Europe. *Pest Manag. Sci.***71**, 492–500.
- Massei, G., Coats, J., Lambert, M.S., Pietravalle, S., Gill, R. & Cowan, D. (2017). Camera traps and activity signs to estimate wild boar density and derive abundance indices. *Pest Manag. Sci.***74**, 853–860.
- Meek, P.D., Ballard, G.-A., Fleming, P.J.S., Schaefer, M., Williams, W. & Falzon, G. (2014). Camera Traps Can Be Heard and Seen by Animals. *PLoS One* **9**.
- Mills, K.J., Patterson, B.R. & Murray, D.L. (2006). Effects of variable sampling frequencies on GPS transmitter efficiency and estimated wolf home range size and movement distance. *Wildl. Soc. Bull.***34**, 1463–1469.
- Morelle, K., Podgórski, T., Prevot, C., Keuling, O., Lehaire, F. & Lejeune, P. (2015). Towards understanding wild boar *Sus scrofa* movement: a synthetic movement ecology approach. *Mamm. Rev.* **45**, 15–29.
- Morellet, N., Gaillard, J.-M., Hewison, A.J.M., Ballon, P., Boscardin, Y., Duncan, P., Klein, F. & Maillard, D. (2007). Indicators of ecological change: new tools for managing populations of large herbivores. *J. Appl. Ecol.***44**, 634–643.
- Musiani, M., Okarma, H. & Jędrzejewski, W. (1998). Speed and actual distances travelled by radiocollared wolves in Białowieża Primeval Forest (Poland). *Acta Theriol. (Warsz.)***43**, 409–416.
- Nakashima, Y., Fukasawa, K. & Samejima, H. (2018). Estimating animal density without individual recognition using information derivable exclusively from camera traps. *J. Appl. Ecol.***55**, 735–744.
- Nichols, J. & Williams, B. (2006). Monitoring for conservation. *Trends Ecol. Evol.***21**, 668–673.
- Pépin, D., Adrados, C., Mann, C. & Janeau, G. (2004). Assessing real daily distance

travelled by ungulates using differential GPS locations. *J. Mammal.***85**, 774–780.

Pfeffer, S.E., Spitzer, R., Allen, A.M., Hofmeester, T.R., Ericsson, G., Widemo, F., Singh, N.J. & Cromsigt, J.P.G.M. (2018). Pictures or pellets? Comparing camera trapping and dung counts as methods for estimating population densities of ungulates. *Remote Sens. Ecol. Conserv.***4**, 173–183.

Prichard, A.K., Yokel, D.A., Rea, C.L., Person, B.T. & Parrett, L.S. (2014). The effect of frequency of telemetry locations on movement-rate calculations in arctic caribou. *Wildl. Soc. Bull.***38**, 78–88.

Podgórski, T., Baś, G., Jędrzejewska, B., Sönnichsen, L., Śnieżko, S., Jędrzejewski, W. & Okarma, H. (2013). Spatiotemporal behavioral plasticity of wild boar ( *Sus scrofa* ) under contrasting conditions of human pressure: primeval forest and metropolitan area. *J. Mammal.* **94**, 109–119.

Podgórski, T., Apollonio, M. & Keuling, O. (2018). Contact rates in wild boar populations: Implications for disease transmission. *J. Wildl. Manage.* **82**, 1210–1218.

Podgórski, T. & Śmietanka, K. (2018). Do wild boar movements drive the spread of African Swine Fever? *Transbound. Emerg. Dis.* **65**, 1588–1596.

Rademaker, M., Meijaard, E., Semiadi, G., Blokland, S., Neilson, E.W. & Rode-Margono, E.J. (2016). First Ecological Study of the Bawean Warty Pig (*Sus blouchi*), One of the Rarest Pigs on Earth. *PLoS One***11**, e0151732.

Reynolds, T.D. & Laundre, J.W. (1990). Time Intervals for Estimating Pronghorn and Coyote Home Ranges and Daily Movements. *J. Wildl. Manage.***54**, 316.

Rovero, F. & Marshall, A.R. (2009). Camera trapping photographic rate as an index of density in forest ungulates. *J. Appl. Ecol.***46**, 1011–1017.

Rovero, F. & Zimmermann, F. (2016). *Camera Trapping for Wildlife Research*. Pelagic Pu.

Rowcliffe, J.M., Field, J., Turvey, S.T. & Carbone, C. (2008). Estimating animal density using camera traps without the need for individual recognition. *J. Appl. Ecol.***45**,

548 1228–1236.

549 Rowcliffe, J.M., Carbone, C., Jansen, P.A., Kays, R. & Kranstauber, B. (2011).  
 550 Quantifying the sensitivity of camera traps: an adapted distance sampling approach.  
 551 *Methods Ecol. Evol.***2**, 464–476.

552 Rowcliffe, J.M., Carbone, C., Kays, R., Kranstauber, B. & Jansen, P.A. (2012). Bias in  
 553 estimating animal travel distance: the effect of sampling frequency. *Methods Ecol.*  
 554 *Evol.***3**, 653–662.

555 Rowcliffe, J.M., Kays, R., Carbone, C. & Jansen, P.A. (2013). Clarifying assumptions  
 556 behind the estimation of animal density from camera trap rates. *J. Wildl. Manage.***77**,  
 557 876.

558 Rowcliffe, J.M., Kays, R., Kranstauber, B., Carbone, C. & Jansen, P.A. (2014).  
 559 Quantifying levels of animal activity using camera trap data. *Methods Ecol. Evol.***5**,  
 560 1170–1179.

561 Rowcliffe, J. M. (2016). activity: Animal Activity Statistics. R package version 1.1.  
 562 <https://CRAN.R-project.org/package=activity>

563 Rowcliffe, J.M., Jansen, P.A., Kays, R., Kranstauber, B. & Carbone, C. (2016). Wildlife  
 564 speed cameras: measuring animal travel speed and day range using camera traps.  
 565 *Remote Sens. Ecol. Conserv.***2**, 84–94.

566 Royle, J.A. & Young, K. V. (2008). A hierarchical model for spatial capture–recapture  
 567 data. *Ecology***89**, 2281–2289.

568 Sennhenn-Reulen, H., Diedhiou, L., Klapproth, M. & Zinner, D. (2017). Estimation of  
 569 baboon daily travel distances by means of point sampling – the magnitude of  
 570 underestimation. *Primate Biol.***4**, 143–151.

571 Spitz, F. & Janeau, G. (1990). Spatial strategies: an attempt to classify daily movements  
 572 of wild boar. *Acta Theriol. (Warsz)*. **35**, 129–149.

573 Steenweg, R., Hebblewhite, M., Kays, R., Ahumada, J., Fisher, J.T., Burton, C.,  
 574 Townsend, S.E., Carbone, C., Rowcliffe, J.M., Whittington, J., Brodie, J., Royle,

575 J.A., Switalski, A., Clevenger, A.P., Heim, N. & Rich, L.N. (2017). Scaling-up  
576 camera traps: monitoring the planet's biodiversity with networks of remote sensors.  
577 *Front. Ecol. Environ.* **15**, 26–34.

578 Tabak, M.A., Norouzzadeh, M.S., Wolfson, D.W., Sweeney, S.J., Vercauteren, K.C.,  
579 Snow, N.P., Halseth, J.M., Di Salvo, P.A., Lewis, J.S., White, M.D., Teton, B.,  
580 Beasley, J.C., Schlichting, P.E., Boughton, R.K., Wight, B., Newkirk, E.S., Ivan, J.S.,  
581 Odell, E.A., Brook, R.K., Lukacs, P.M., Moeller, A.K., Mandeville, E.G., Clune, J. &  
582 Miller, R.S. (2018). Machine learning to classify animal species in camera trap  
583 images: Applications in ecology. *Methods Ecol. Evol.*

584 Tucker, M.A., Böhning-Gaese, K., Fagan, W.F., Fryxell, J.M., Van Moorter, B., Alberts,  
585 S.C., Ali, A.H., Allen, A.M., Attias, N., Avgar, T., Bartlam-Brooks, H., Bayarbaatar,  
586 B., Belant, J.L., Bertassoni, A., Beyer, D., Bidner, L., van Beest, F.M., Blake, S.,  
587 Blaum, N., Bracis, C., Brown, D., de Bruyn, P.J.N., Cagnacci, F., Calabrese, J.M.,  
588 Camilo-Alves, C., Chamaillé-Jammes, S., Chiaradia, A., Davidson, S.C., Dennis,  
589 T., DeStefano, S., Diefenbach, D., Douglas-Hamilton, I., Fennessy, J., Fichtel, C.,  
590 Fiedler, W., Fischer, C., Fischhoff, I., Fleming, C.H., Ford, A.T., Fritz, S.A., Gehr,  
591 B., Goheen, J.R., Gurarie, E., Hebblewhite, M., Heurich, M., Hewison, A.J.M., Hof,  
592 C., Hurme, E., Isbell, L.A., Janssen, R., Jeltsch, F., Kaczensky, P., Kane, A.,  
593 Kappeler, P.M., Kauffman, M., Kays, R., Kimuyu, D., Koch, F., Kranstauber, B.,  
594 LaPoint, S., Leimgruber, P., Linnell, J.D.C., López-López, P., Markham, A.C.,  
595 Mattisson, J., Medici, E.P., Mellone, U., Merrill, E., de Miranda Mourão, G., Morato,  
596 R.G., Morellet, N., Morrison, T.A., Díaz-Muñoz, S.L., Mysterud, A., Nandintsetseg,  
597 D., Nathan, R., Niamir, A., Odden, J., O'Hara, R.B., Oliveira-Santos, L.G.R., Olson,  
598 K.A., Patterson, B.D., Cunha de Paula, R., Pedrotti, L., Reineking, B., Rimmler, M.,  
599 Rogers, T.L., Rolandsen, C.M., Rosenberry, C.S., Rubenstein, D.I., Safi, K., Saïd,  
600 S., Sapir, N., Sawyer, H., Schmidt, N.M., Selva, N., Sergiel, A., Shiilegdamba, E.,  
601 Silva, J.P., Singh, N., Solberg, E.J., Spiegel, O., Strand, O., Sundaresan, S.,  
602 Ullmann, W., Voigt, U., Wall, J., Wattles, D., Wikelski, M., Wilmers, C.C., Wilson,

J.W., Wittemyer, G., Zięba, F., Zwijacz-Kozica, T. & Mueller, T. (2018). Moving in the Anthropocene: Global reductions in terrestrial mammalian movements. *Science* (80-. ).**359**, 466–469.

Wald, A. & Wolfowitz, J. (1940). On a Test Whether Two Samples are from the Same Population. *Ann. Math. Stat.***11**, 147–162.


Zero, V.H., Sundaresan, S.R., O'Brien, T.G. & Kinnaird, M.F. (2013). Monitoring an Endangered savannah ungulate, Grevy's zebra *Equus grevyi*: choosing a method for estimating population densities. *Oryx*.**47**, 410–419.

**Table 1.** Summary of movement parameters estimated using camera-traps. Travel speed estimations (average speed of travel while active) are based on the best-fit distribution selected on the basis of AIC. Activity index estimates (proportion of time spent active) were calculated using the R package “activity”


Camera mode	Approach	Activity index (SE)	Travel speed m·s <sup>-1</sup> (SE)	Speed distribution		Day range Km·day <sup>-1</sup> (SE)
				(ΔAIC)		
				Log-normal	Weibull	
Photo	General	0.361	0.158	6.76	0	4.928
		(0.034)	(0.029)			(1.020)
	Weighted	0.150 <sup>f</sup>	0.033 <sup>f</sup>	3.01 <sup>f</sup>	0 <sup>f</sup>	12.740
		(0.014)	(0.016)			(2.073)
Video	General	0.211 <sup>m</sup>	0.632 <sup>m</sup>	6.86 <sup>m</sup>	0 <sup>m</sup>	3.014
		(0.020)	(0.075)			(1.064)
	Weighted	0.171 <sup>f</sup>	0.072 <sup>f</sup>	0 <sup>f</sup>	1.60 <sup>f</sup>	13.997
		(0.017)	(0.019)			(2.404)
		0.221 <sup>m</sup>	0.680 <sup>m</sup>	0 <sup>m</sup>	3.21 <sup>m</sup>	
		(0.022)	(0.088)			

<sup>f</sup> feeding behaviour, <sup>m</sup> moving behaviour.







**Figure 1.** Map of the study area. Camera-traps locations (black points) and home range of tagged individuals in period 3 (dashed line).



**Figure 2.** A) Scheme of the stick-structure (grey dots) used to reference the animals captured by the camera-trap (black dot).  $X_c$  and  $X_d$  indicate the position of the wild boar captured in images C and D, respectively. B) Photo of the structure installed at one photo-trapping sampling point. C) & D) Photographic sequence of a wild boar captured by the camera.



**Figure 3.** Activity patterns of wild boar estimated from photo (A) and video data (B) using R package “activity”. Black steps are observed frequencies, grey curves are fitted kernel distributions and grey discontinuous lines are SE.



**Figure 4.** Day range estimates obtained from: approach-1.1 to approach-1.3, telemetry data corrected by tortuosity-related correction factor (Rowcliffe *et al.*, 2012) corresponding with the periods 1, 2 and 3 respectively; approach-2.1, general estimation based on camera-trap photos; approach-2.2, general estimation based on camera-trap videos; approach-3.1, weighted estimation based on camera-trap photos; approach-3.2, weighted estimation based on camera-trap videos. Upper axis indicates time lapse between consecutive locations (in mins) in telemetry data. Lower black bars in telemetry estimates correspond to straight-line estimates and upper white bars are the tortuosity-related correction in approaches 1.1-1.3. Different letters show significant differences among DR estimation based on pairwise comparisons with Bonferroni correction ( $p < 0.05$ ) after Wald test analysis. Error bars are 95% confidence intervals.